


CANCER/ANTI-CANCER DUAL ACTION DRUGS IN CROWN-GALL TUMORS

L. Le Goff and M. Beljanski

Laboratoire de Pharmacodynamie, Faculté de Pharmacie, Rue J.B. Clement, 92290, Châtenay-Malabry, France

Paper received: 2nd July, 1979; amended 29th August, 1979

Parallel studies on plant and animal tumor systems provide insight into the basic cellular mechanisms of tumorigenesis. In laboratory animals, drugs characterized as mutagens with the *Salmonella* assay system (1), as for example cyclophosphamide, may have an antimitotic effect and/or give rise to neoplasms (2). Various other *in vitro* carcinogen screening tests



Abcissae: drug concentration expressed as μg per wound (Log C). Ordinate: average of the adjusted tumor weight expressed as percentage of control (W). Drugs were applied to the wounds either mixed with *B₆* bacteria (10^8 cells/wound) (T_0 : $\bullet-\bullet-\bullet$) or separately, 24 h ($\circ-\circ-\circ$) or 48 h ($\Delta-\Delta-\Delta$) after bacterial infection. Each point represents mean tumor weight value for 50 plants, and means \pm SE are shown.

day, tumors were excised and weighed and lengths of lateral shoots were measured. As the tumor weight increases, the length of shoots decreases. We calculated as described (5) for each plant an estimated tumor weight corresponding to non-existent lateral shoots. Fifty plants were used for each assay. Means of adjusted values were statistically compared with a confidence interval of 1%, according to Student's *t* test.

Results: The figure shows that, at low concentrations, cyclophosphamide and Daunorubicin act as tumor-stimulants when introduced into wounded tissues as early as 24 or 48 h following infection with *A. tumefaciens* (a 24 h lapse is needed for conversion of healthy plant cells into tumor cells). Four days after inoculation of *B₆* strain (this is the time period during which the bacteria are no longer tumor-inducing) stimulation of neoplastic cell growth by both drugs can still be observed. At the same low concentrations, these drugs have no effect when introduced together with the bacteria at time 0, that is to say during induction of cancer tissue formation. At high concentrations, cyclophosphamide and Daunorubicin, which have apparently no damaging effect on plant growth (lateral shoot length measurements remain normal during the entire 12 days experimental period), show a strong tumor inhibiting action (figure). It must be emphasized that in the absence of bacteria neither of these drugs has any detectable carcinogenic effect under the experimental conditions we use.

Conclusions: Carcinogenic or/and anti-cancer potency of some therapeutic drugs may be established *in vivo* using plant material which is easy to handle, is not costly and produces results in a relatively short period of time (12 days). In animals the effect of these drugs requires a very long period of time to be detected. Our *in vivo* observations of the carcinogenic and anti-cancer potency of cyclophosphamide and Daunorubicin are in agreement with data obtained with these same drugs in *in vitro* tests or observed in animals (2) and humans (6), and point a possible recognition of their effect in plant systems.

1. Benedict, W.F., et al., (1977) *Cancer Res.*, 37, 2209
2. Schmähl, D. and Habs, M. (1979) *Int. J. Cancer*, 23, 706
3. Lenzin, A., et al., (1978) *Urologe A*, 17, 105
4. Beljanski, M. (1979) *IRCS Med. Sci.*, 7, 476
5. Kurkdjian, A., et al., (1974) *Am. Biol. Teacher*, 36, 13
6. Bertazzoli, G., et al., (1971) *Experientia*, 27, 1209


CANCER/ANTI-CANCER DUAL ACTION DRUGS IN CROWN-GALL TUMORS

L. Le Goff and M. Beljanski

Laboratoire de Pharmacodynamie, Faculté de Pharmacie, Rue J.B. Clement, 92290, Châtenay-Malabry, France

Paper received: 2nd July, 1979; amended 29th August, 1979

Parallel studies on plant and animal tumor systems provide insight into the basic cellular mechanisms of tumorigenesis. In laboratory animals, drugs characterized as mutagens with the *Salmonella* assay system (1), as for example cyclophosphamide, may have an antimitotic effect and/or give rise to neoplasms (2). Various other *in vitro* carcinogen screening tests



Abscissae: drug concentration expressed as μg per wound (Log C). Ordinates: average of the adjusted tumor weight expressed as percentage of control (W). Drugs were applied to the wounds either mixed with *B₆* bacteria (10^9 cells/wound) (T_0 : ●—●—●) or separately, 24 h (○—○—○) or 48 h (▲...▲...▲) after bacterial infection. Each point represents mean tumor weight value for 50 plants, and means \pm SE are shown.

day, tumors were excised and weighed and lengths of lateral shoots were measured. As the tumor weight increases, the length of shoots decreases. We calculated as described (5) for each plant an estimated tumor weight corresponding to non-existent lateral shoots. Fifty plants were used for each assay. Means of adjusted values were statistically compared with a confidence interval of 1%, according to Student's *t* test.

Results: The figure shows that, at low concentrations, cyclophosphamide and Daunorubicin act as tumor-stimulants when introduced into wounded tissues as early as 24 or 48 h following infection with *A. tumefaciens* (a 24 h lapse is needed for conversion of healthy plant cells into tumor cells). Four days after inoculation of *B₆* strain (this is the time period during which the bacteria are no longer tumor-inducing) stimulation of neoplastic cell growth by both drugs can still be observed. At the same low concentrations, these drugs have no effect when introduced together with the bacteria at time 0, that is to say during induction of cancer tissue formation. At high concentrations, cyclophosphamide and Daunorubicin, which have apparently no damaging effect on plant growth (lateral shoot length measurements remain normal during the entire 12 days experimental period), show a strong tumor inhibiting action (figure). It must be emphasized that in the absence of bacteria neither of these drugs has any detectable carcinogenic effect under the experimental conditions we use.

Conclusions: Carcinogenic or/and anti-cancer potency of some therapeutic drugs may be established *in vivo* using plant material which is easy to handle, is not costly and produces results in a relatively short period of time (12 days). In animals the effect of these drugs requires a very long period of time to be detected. Our *in vivo* observations of the carcinogenic and anti-cancer potency of cyclophosphamide and Daunorubicin are in agreement with data obtained with these same drugs in *in vitro* tests or observed in animals (2) and humans (6), and point a possible recognition of their effect in plant systems.

have shown that cyclophosphamide and Daunorubicin have mutagenic (1) and carcinogenic potencies (3, 4). We give here a brief summary of the effect of two anticancer drugs, cyclophosphamide (Endoxan) and Danorubicin, on the heritable cellular change in pre-induced Crown-gall tumor development.

Material and methods: Germinating decapitated pea seedlings (*Pisum sativum* c.v. Annonay (Vilmorin)) experimentally infected with *Agrobacterium tumefaciens*, tumorigenic strain *B₆*, were used as already described (5). Drugs were dissolved in buffered physiological saline solution (pH 7.0) and filtered on a millipore filter before use. They were introduced into the wound either mixed with bacteria (time 0) or separately, by tissue scarification, at various times after start of bacterial infection (figure). On the 12th

1. Benedict, W.F., et al., (1977) *Cancer Res.*, 37, 2209
2. Schmähl, D. and Habs, M. (1979) *Int. J. Cancer*, 23, 706
3. Lenzin, A., et al., (1978) *Urologe A*, 17, 105
4. Beljanski, M. (1979) *IRCS Med. Sci.*, 7, 476
5. Kurkdjian, A., et al., (1974) *Am. Biol. Teacher*, 36, 13
6. Bertazzoli, G., et al., (1971) *Experientia*, 27, 1209


CANCER/ANTI-CANCER DUAL ACTION DRUGS IN CROWN-GALL TUMORS

L. Le Goff and M. Beljanski

Laboratoire de Pharmacodynamie, Faculté de Pharmacie, Rue J.B. Clement, 92290, Châtenay-Malabry, France

Paper received: 2nd July, 1979; amended 29th August, 1979

Parallel studies on plant and animal tumor systems provide insight into the basic cellular mechanisms of tumorigenesis. In laboratory animals, drugs characterized as mutagens with the *Salmonella* assay system (1), as for example cyclophosphamide, may have an antimitotic effect and/or give rise to neoplasms (2). Various other *in vitro* carcinogen screening tests



Abscissae: drug concentration expressed as μg per wound (Log C). Ordinates: average of the adjusted tumor weight expressed as percentage of control (W). Drugs were applied to the wounds either mixed with *B₆* bacteria (10^8 cells/wound) (T_0 : ●—●) or separately, 24 h (○—○) or 48 h (▲...▲) after bacterial infection. Each point represents mean tumor weight value for 50 plants, and means \pm SE are shown.

day, tumors were excised and weighed and lengths of lateral shoots were measured. As the tumor weight increases, the length of shoots decreases. We calculated as described (5) for each plant an estimated tumor weight corresponding to non-existent lateral shoots. Fifty plants were used for each assay. Means of adjusted values were statistically compared with a confidence interval of 1%, according to Student's *t* test.

Results: The figure shows that, at low concentrations, cyclophosphamide and Daunorubicin act as tumor-stimulants when introduced into wounded tissues as early as 24 or 48 h following infection with *A. tumefaciens* (a 24 h lapse is needed for conversion of healthy plant cells into tumor cells). Four days after inoculation of *B₆* strain (this is the time period during which the bacteria are no longer tumor-inducing) stimulation of neoplastic cell growth by both drugs can still be observed. At the same low concentrations, these drugs have no effect when introduced together with the bacteria at time 0, that is to say during induction of cancer tissue formation. At high concentrations, cyclophosphamide and Daunorubicin, which have apparently no damaging effect on plant growth (lateral shoot length measurements remain normal during the entire 12 days experimental period), show a strong tumor inhibiting action (figure). It must be emphasized that in the absence of bacteria neither of these drugs has any detectable carcinogenic effect under the experimental conditions we use.

Conclusions: Carcinogenic or/and anti-cancer potency of some therapeutic drugs may be established *in vivo* using plant material which is easy to handle, is not costly and produces results in a relatively short period of time (12 days). In animals the effect of these drugs requires a very long period of time to be detected. Our *in vivo* observations of the carcinogenic and anti-cancer potency of cyclophosphamide and Daunorubicin are in agreement with data obtained with these same drugs in *in vitro* tests or observed in animals (2) and humans (6), and point a possible recognition of their effect in plant systems.

1. Benedict, W.F., et al., (1977) *Cancer Res.*, 37, 2209
2. Schmähl, D. and Habs, M. (1979) *Int. J. Cancer*, 23, 706
3. Lenzin, A., et al., (1978) *Urologe A*, 17, 105
4. Beljanski, M. (1979) *IRCS Med. Sci.*, 7, 476
5. Kurkdjian, A., et al., (1974) *Am. Biol. Teacher*, 36, 13
6. Bertazzoli, G., et al., (1971) *Experientia*, 27, 1209


CANCER/ANTI-CANCER DUAL ACTION DRUGS IN CROWN-GALL TUMORS

L. Le Goff and M. Beljanski

Laboratoire de Pharmacodynamie, Faculté de Pharmacie, Rue J.B. Clement, 92290, Châtenay-Malabry, France

Paper received: 2nd July, 1979; amended 29th August, 1979

Parallel studies on plant and animal tumor systems provide insight into the basic cellular mechanisms of tumorigenesis. In laboratory animals, drugs characterized as mutagens with the *Salmonella* assay system (1), as for example cyclophosphamide, may have an antimitotic effect and/or give rise to neoplasms (2). Various other *in vitro* carcinogen screening tests



Abscissae: drug concentration expressed as μg per wound (Log C). Ordinates: average of the adjusted tumor weight expressed as percentage of control (\bar{W}). Drugs were applied to the wounds either mixed with *B₆* bacteria (10^6 cells/wound) (T_0 : \bullet - \bullet - \bullet) or separately, 24 h (\circ - \circ - \circ) or 48 h (Δ - Δ - Δ) after bacterial infection. Each point represents mean tumor weight value for 50 plants, and means \pm SE are shown.

day, tumors were excised and weighed and lengths of lateral shoots were measured. As the tumor weight increases, the length of shoots decreases. We calculated as described (5) for each plant an estimated tumor weight corresponding to non-existent lateral shoots. Fifty plants were used for each assay. Means of adjusted values were statistically compared with a confidence interval of 1%, according to Student's *t* test.

Results: The figure shows that, at low concentrations, cyclophosphamide and Daunorubicin act as tumor-stimulants when introduced into wounded tissues as early as 24 or 48 h following infection with *A. tumefaciens* (a 24 h lapse is needed for conversion of healthy plant cells into tumor cells). Four days after inoculation of *B₆* strain (this is the time period during which the bacteria are no longer tumor-inducing) stimulation of neoplastic cell growth by both drugs can still be observed. At the same low concentrations, these drugs have no effect when introduced together with the bacteria at time 0, that is to say during induction of cancer tissue formation. At high concentrations, cyclophosphamide and Daunorubicin, which have apparently no damaging effect on plant growth (lateral shoot length measurements remain normal during the entire 12 days experimental period), show a strong tumor inhibiting action (figure). It must be emphasized that in the absence of bacteria neither of these drugs has any detectable carcinogenic effect under the experimental conditions we use.

Conclusions: Carcinogenic or/and anti-cancer potency of some therapeutic drugs may be established *in vivo* using plant material which is easy to handle, is not costly and produces results in a relatively short period of time (12 days). In animals the effect of these drugs requires a very long period of time to be detected. Our *in vivo* observations of the carcinogenic and anti-cancer potency of cyclophosphamide and Daunorubicin are in agreement with data obtained with these same drugs in *in vitro* tests or observed in animals (2) and humans (6), and point a possible recognition of their effect in plant systems.

1. Benedict, W.F., et al., (1977) *Cancer Res.*, 37, 2209
2. Schmähl, D. and Habs, M. (1979) *Int. J. Cancer*, 23, 706
3. Lenzin, A., et al., (1978) *Urologe A*, 17, 105
4. Beljanski, M. (1979) *IRCS Med. Sci.*, 7, 476
5. Kurkdjian, A., et al., (1974) *Am. Biol. Teacher*, 36, 13
6. Bertazzoli, G., et al., (1971) *Experientia*, 27, 1209

have shown that cyclophosphamide and Daunorubicin have mutagenic (1) and carcinogenic potencies (3, 4). We give here a brief summary of the effect of two anticancer drugs, cyclophosphamide (Endoxan) and Danorubicin, on the heritable cellular change in pre-induced Crown-gall tumor development.

Material and methods: Germinating decapitated pea seedlings (*Pisum sativum* c.v. Annonay (Vilmorin)) experimentally infected with *Agrobacterium tumefaciens*, tumorigenic strain *B₆*, were used as already described (5). Drugs were dissolved in buffered physiological saline solution (pH 7.0) and filtered on a millipore filter before use. They were introduced into the wound either mixed with bacteria (time 0) or separately, by tissue scarification, at various times after start of bacterial infection (figure). On the 12th


CANCER/ANTI-CANCER DUAL ACTION DRUGS IN CROWN-GALL TUMORS

L. Le Goff and M. Beljanski

Laboratoire de Pharmacodynamie, Faculté de Pharmacie, Rue J.B. Clement, 92290, Châtenay-Malabry, France

Paper received: 2nd July, 1979; amended 29th August, 1979

Parallel studies on plant and animal tumor systems provide insight into the basic cellular mechanisms of tumorigenesis. In laboratory animals, drugs characterized as mutagens with the *Salmonella* assay system (1), as for example cyclophosphamide, may have an antimutagenic effect and/or give rise to neoplasms (2). Various other *in vitro* carcinogen screening tests



Abscissae: drug concentration expressed as μg per wound (Log C). Ordinates: average of the adjusted tumor weight expressed as percentage of control (W). Drugs were applied to the wounds either mixed with *B₆* bacteria (10^8 cells/wound) (T_0 : $\bullet-\bullet-\bullet$) or separately, 24 h ($\circ-\circ-\circ$) or 48 h ($\Delta-\Delta-\Delta$) after bacterial infection. Each point represents mean tumor weight value for 50 plants, and means \pm SE are shown.

day, tumors were excised and weighed and lengths of lateral shoots were measured. As the tumor weight increases, the length of shoots decreases. We calculated as described (5) for each plant an estimated tumor weight corresponding to non-existent lateral shoots. Fifty plants were used for each assay. Means of adjusted values were statistically compared with a confidence interval of 1%, according to Student's *t* test.

Results: The figure shows that, at low concentrations, cyclophosphamide and Daunorubicin act as tumor-stimulants when introduced into wounded tissues as early as 24 or 48 h following infection with *A. tumefaciens* (a 24 h lapse is needed for conversion of healthy plant cells into tumor cells). Four days after inoculation of *B₆* strain (this is the time period during which the bacteria are no longer tumor-inducing) stimulation of neoplastic cell growth by both drugs can still be observed. At the same low concentrations, these drugs have no effect when introduced together with the bacteria at time 0, that is to say during induction of cancer tissue formation. At high concentrations, cyclophosphamide and Daunorubicin, which have apparently no damaging effect on plant growth (lateral shoot length measurements remain normal during the entire 12 days experimental period), show a strong tumor inhibiting action (figure). It must be emphasized that in the absence of bacteria neither of these drugs has any detectable carcinogenic effect under the experimental conditions we use.

Conclusions: Carcinogenic or/and anti-cancer potency of some therapeutic drugs may be established *in vivo* using plant material which is easy to handle, is not costly and produces results in a relatively short period of time (12 days). In animals the effect of these drugs requires a very long period of time to be detected. Our *in vivo* observations of the carcinogenic and anti-cancer potency of cyclophosphamide and Daunorubicin are in agreement with data obtained with these same drugs in *in vitro* tests or observed in animals (2) and humans (6), and point a possible recognition of their effect in plant systems.

1. Benedict, W.F., et al., (1977) *Cancer Res.*, 37, 2209
2. Schmähl, D. and Habs, M. (1979) *Int. J. Cancer*, 23, 706
3. Lenzin, A., et al., (1978) *Urologe A*, 17, 105
4. Beljanski, M. (1979) *IRCS Med. Sci.*, 7, 476
5. Kurkdjian, A., et al., (1974) *Am. Biol. Teacher*, 36, 13
6. Bertazzoli, G., et al., (1971) *Experientia*, 27, 1209

have shown that cyclophosphamide and Daunorubicin have mutagenic (1) and carcinogenic potencies (3, 4). We give here a brief summary of the effect of two anticancer drugs, cyclophosphamide (Endoxan) and Daunorubicin, on the heritable cellular change in pre-induced Crown-gall tumor development.

Material and methods: Germinating decapitated pea seedlings (*Pisum sativum* c.v. Annonay (Vilmorin)) experimentally infected with *Agrobacterium tumefaciens*, tumorigenic strain *B₆*, were used as already described (5). Drugs were dissolved in buffered physiological saline solution (pH 7.0) and filtered on a millipore filter before use. They were introduced into the wound either mixed with bacteria (time 0) or separately, by tissue scarification, at various times after start of bacterial infection (figure). On the 12th